arXiv:1410.5181v1 [math.MG] 20 Oct 2014

ON STATIC EQUILIBRIA AND ISOPERIMETRIC RATIO OF AFFINE FAMILIES OF CONVEX BODIES

GÁBOR DOMOKOS AND ZSOLT LÁNGI

Abstract. Motivated by physical abrasion processes, in this note we prove some simple facts about the evolution of the isoperimetric ratio of convex bodies subject to one-parameter orthogonal affinity. We also show how the number of static equilibrium points is evolving if the parameter of the affinity tends to zero or to infinity.

1. Introduction

Physical abrasion processes are composed of collisional and frictional abrasion [3],[4]. While the former is relatively well understood ([2], [7]), little is known about the latter. The basic geometric axioms for frictional abrasion are defined in [3], however, many particular evolution models satisfy those axioms and it is not yet clear which of these is physically relevant. Recently, new physical indicators of geological abrasion processes have been investigated, including the evolution of the isoperimetric ratio I, which has been measured both in the field [9] and in laboratory experiments [10]. The isoperimetric ratio I is of particular interest, because in case of curvature-driven flows (serving as models of *collisional* abrasion) it was proven for planar curves by Gage [8] that $dI/dt > 0$. In this note we postulate one-parameter (t) orthogonal affinity as one particular model of frictional abrasion (satisfying the axioms in [3]) and we describe some aspects of the evolution of the isoperimetric ratio. In particular, we show that $I(t)$ is a quasiconcave function (Theorem 1) and we also show that there exists an orthogonal basis along the directions of which $dI/dt = 0$ (Theorem 2). Our results hold in arbitrary dimensions.

Another, recently investigated indicator of abrasion processes is the number T of static equilibrium points [5]. In two dimensions, (planar disks) we distinguish between stable and unstable equilibria, and denote their numbers respectively by S and U and we have $S + U = T$. In three dimensions (solids) we distinguish between stable, unstable and saddle-type equilibria, and denote their numbers respectively by S, U, N and on generic (Morse) surfaces we have $S+U+N = T$. These numbers are related by the the Poincaré-Hopf Theorem [1] on topological invariants as

$$S - U = 0 \text{ (in 2 dimensions)}, \quad S + U - N = 2 \text{ (in 3 dimensions)}. \tag{1}$$

The evolution of these numbers has been already measured in the field [9]. In the current note we will study their evolution under orthogonal affinity. In particular we will show (Theorem 3) that in two dimensions, as t approaches either zero or infinity, $T(t)$ approaches 4. In 3 dimensions, as t approaches zero or infinity, either $S(t)$ or $U(t)$ approach 2. We also formulate additional statements about these numbers and a conjecture about the global connection between $I(t)$ and $T(t)$ in an averaged sense.

1991 *Mathematics Subject Classification.* 53A05, 53Z05,52A38.

Key words and phrases. equilibrium, convex surface, affinity, isoperimetric ratio.

The authors gratefully acknowledge the support of the János Bolyai Research Scholarship of the Hungarian Academy of Sciences and support from OTKA grant 104601.

2. Main results

Let $K \subset \mathbb{R}^n$ be a convex body with C^2-class boundary and nowhere vanishing curvature, and H be a hyperplane passing through the centre of gravity of K, with normal vector $v \in \mathbb{S}^{n-1}$. Following [6], we call the quantity $\frac{\mathrm{vol}(K)}{(\mathrm{surf}(K))^{\frac{n}{n-1}}}$ the *isoperimetric ratio of* K.

Let $a_t^v : \mathbb{R}^n \to \mathbb{R}^n$ be the orthogonal affinity, with H as its fixed hyperplane, and ratio $t > 0$, and set $K^v(t) = a_t^v(K)$.

Theorem 1. *For every convex body $K \subseteq \mathbb{R}^n$ and every $v \in \mathbb{S}^{n-1}$, the isoperimetric ratio of $K^v(t)$ is a quasiconcave function of t.*

Proof. Set $V(t) = \mathrm{vol}\, K^v(t)$ and $A(t) = \mathrm{surf}\, K^v(t)$. Without loss of generality, we may assume that $V(1) = 1$ which implies that $V(t) = t$. Thus we need to show that the function $I(t) = \frac{t}{A^s(t)}$, where $s = \frac{n}{n-1}$, is quasiconcave. We show that $I'(t)$ has exactly one root, which, since $I(t) \to 0$ as $t \to 0$ or $t \to \infty$, yields that here $I(t)$ has a maximum.

Observe that $I'(t) = \frac{A(t) - stA'(t)}{A^{s+1}(t)}$, and that $(A(t) - stA'(t))' = (1-s)A'(t) - stA''(t)$. Note that as $1 - s < 0$ and $A' > 0$, if $A''(t) < 0$, then the numerator of $I'(t)$ is strictly decreasing, which, combined with the limits $I(t) \to 0$ as $t \to 0$ or $t \to \infty$, implies the assertion.

We show that $A''(t) < 0$. Let us imagine H as the hyperplane $\{x_n = 0\}$, and let D denote the orthogonal projection of K onto H. Then $\mathrm{bd}\, K$ is the union of the graphs of two functions defined on D, and thus, it is enough to prove the statement for graphs of functions.

Consider the function $x_n = f(x_1, x_2, \ldots, x_{n-1})$ defined on D. Then, applying the orthogonal affinity with ratio t we obtain the function $x_n = tf(x_1, x_2, \ldots, x_{n-1})$, the surface area of which we denote by $A_f(t)$. Then

$$A_f''(t) = \frac{\mathrm{d}^2}{\mathrm{d}\, t^2} \int_{x \in D} \sqrt{1 + t^2 \,\mathrm{grad}^2\, f}\, \mathrm{d}\, x = \int_{x \in D} \frac{\mathrm{d}^2}{\mathrm{d}\, t^2} \sqrt{1 + t^2 \,\mathrm{grad}^2\, f}\, \mathrm{d}\, x =$$

$$= \int_{x \in D} \frac{\mathrm{grad}^2\, f}{\left(1 + t^2 \,\mathrm{grad}^2\, f\right)^{3/2}}\, \mathrm{d}\, x,$$

where the integrand is positive. □

Theorem 2. *Let $K \subset \mathbb{R}^n$ be a convex body. For any $v \in \mathbb{S}^{n-1}$, let $I_v(t)$ denote the isoperimetric ratio of $a^v(K(t))$. Then there is an orthonormal basis $e_1, e_2, \ldots, e_n \in \mathbb{R}^n$ such that for any i, $I'_{e_i}(1) = 0$.*

Proof. For any $v \in \mathbb{S}^{n-1}$ let $f(v) = I'_v(1)$. Then, since $\mathrm{bd}\, K$ is C^2-class differentiable, $f : \mathbb{S}^{n-1} : \mathbb{R}$ is continuous. We show that for any orthonormal basis $e_1, e_2, \ldots, e_n$, we have $\sum_{i=1}^n f(e_i) = 0$.

Given an orthonormal basis, we define a function $g : \mathbb{R}^n \to \mathbb{R}$ in the following way: For $i = 1, 2, \ldots, n$, let H_i be a hyperplane orthogonal to e_i, and a_t^i the orthogonal affinity, of ratio t, with H_i as its fixed hyperplane. We set $K(t_1, t_2, \ldots, t_n) = a^1_{t_1}(a^2_{t_2}(\ldots a^n_{t_n}(K)\ldots))$, and note that this body is independent of the order in which the affinities are carried out. Finally, let $g(t_1, t_2 \ldots, t_n)$ be the isoperimetric ratio of the body $K(t_1, t_2, \ldots, t_n)$. Clearly, this function is differentiable at $(1, 1, \ldots, 1)$. Let $v = \sum_{i=1}^n e_i$. Then, by the linearity of the directional derivatives, we have $g'_v(1, \ldots, 1) = \sum_{i=1}^n g'_{e_i}(1, \ldots, 1) = \sum_{i=1}^n f(e_i)$. On

the other hand, $g(t,\dots,t) = g(1,\dots,1)$ clearly holds for every $t > 0$, which yields that $g'_v(1,\dots,1) = 0$.

Now we prove the following, more general statement: If $h : \mathbb{S}^{n-1} : \mathbb{R}$ is a continuous function such that for every orthonormal basis $e_1, e_2, \dots, e_n$, we have $\sum_{i=1}^n h(e_i) = 0$, then there is an orthonormal basis $e_1, e_2, \dots, e_n$ such that $h(e_i) = 0$ for every value of i. This clearly implies the assertion.

We show this statement by induction on n. If $n = 1$, then the statement is obvious. Now, assume that the statement holds for functions defined on $\mathbb{S}^{n-2}$. Consider some $h : \mathbb{S}^{n-1} : \mathbb{R}$ satisfying our conditions. Then there are some (orthogonal) vectors $u, w \in \mathbb{S}^{n-1}$ such that $h(u) \le 0 \le h(w)$. Thus, there is some $v \in \mathbb{S}n-1$ such that $h(v) = 0$. We identify the set of vectors in $\mathbb{R}^n$, perpendicular to v with the space $\mathbb{R}^{n-1}$, and let h_v denote restriction of h to this subspace. Then, if $e_1, \dots, e_{n-1} \in \mathbb{S}^{n-2} = \mathbb{R}^{n-1} \cap \mathbb{S}^{n-1}$ is an orthogonal basis in $\mathbb{R}^{n-1}$, then, adding v to it we obtain an orthogonal basis in $\mathbb{R}^n$, and thus, we obtain that $\sum_{i=1}^{n-1} h(e_i) = h(v) + \sum_{i=1}^{n-1} h(e_i) = 0$. Hence, we can apply the inductive hypothesis to h_v, which yields the required statement. □

Note that we have proven the following, stronger statement.

Corollary 1. *Any orthonormal k-frame $e_1, e_2, \dots, e_k$ in $\mathbb{R}^n$, satisfying $I'_{e_i}(1) = 0$ for every value of i, can be completed to an orthonormal basis $e_1, \dots, e_n$ satisfying the same property.*

Theorem 3. *Let $u_K(t)$ denote the number of unstable points of $K(t)$, with respect to its centre of gravity. Then, if t is sufficiently large, then $u_K(t) = 2$.*

Proof. First, we prove the assertion for $n = 2$.

Let the origin o be the centre of gravity of K, and H the x-axis. Let the projection of K onto the x axis be $[a, b]$, with $a < 0$ and $b > 0$. Then $\operatorname{bd} K$ is the union of the graphs of two functions, defined on $[a, b]$, one strictly concave and the other one strictly convex. Let $f : [a, b] \to \mathbb{R}$ be the strictly concave one, which is then C^2-class, and has nonvanishing curvature. Then $\operatorname{bd} K(t)$ can be written as the set $\{(x, tf(x)) \in \mathbb{R}^2 : x \in [a, b]$.

Let $x_0 \in (a, b)$ be the value with $f'(x_0) = 0$, and note that x_0 uniquely exists, as under our conditions, f' is strictly decreasing. First, observe that $K(t)$ has an equilibrium at $(x, tf(x))$ if, and only if the tangent line of f at this point is perpendicular to the position vector of the point, or in other words, if

$$0 = \langle (x, tf(x)), (1, tf'(x)) \rangle = x + t^2 f(x) f'(x). \tag{2}$$

For any fixed x with $f'(x) \neq 0 \neq f(x)$, the right-hand side expression is a strictly monotonous function of t, which means that it is satisfied for exactly one value of t. Thus for any $\varepsilon > 0$, if t is sufficiently large, then we have one of the following for any equilibrium point $(x, tf(x))$ of $K(t)$ on f:

(1) $|x - x_0| < \varepsilon$;
(2) $|x - x'_0| < \varepsilon$ for some x'_0 satisfying $f(x'_0) = 0$.

First, we consider the first type equilibria. Note that $f(x_0) > 0$, $f'(x_0)$ and by concavity and the nonvanishing of curvature, $f''(x_0) < 0$. Thus, there is some value t_0 such that $t_0^2 \left((f'(x_0))^2 + f(x_0) f''(x_0) \right) < -1$. Furthermore, since $f(x)f''(x) + (f'(x))^2$ is a continuous function of x, there is some neighborhood U of x_0 such that $t_0^2 \left((f'(x))^2 + f(x) f''(x) \right) < -1$ holds for any $x \in U$. Note that for any $x \in U$, the same inequality holds for any $t > t_0$.

As the x-derivative of the right-hand side $R_t(x)$ of (2) is $1 + t^2 \left((f'(x))^2 + f(x) f''(x) \right)$, we obtain that for any $t \ge t_0$, $R_t(x)$ is a strictly decreasing function on U. Hence, choosing

$\varepsilon > 0$ such that $(x_0 - \varepsilon, x_0 + \varepsilon) \subset U$, for every sufficiently large $t \geq t_0$, the (2) is satisfied for exactly one value x_t of $x \in (a, b)$. Furthermore, in this case for any $x \in U$, $x < x_t$ yields $R_t(x) > 0$ and $x > x_t$ yields $R_t(x) < 0$, it follows that tK has an unstable point at $(x_t, tf(x_t))$.

Now we consider a second type equilibrium point x_t'. Applying a similar argument as in the previous case, one can see that if t is sufficiently large, then, in a neighborhood of x_0', the Euclidean distance function of $\operatorname{bd} K$ is minimal at x_t, which yields that t is a stable point of K. Thus, if t is sufficiently large, then $u_K(t) = 2$.

Now, we prove the statement for any dimension $n > 2$. Let $H = \{x_n = 0\}$, which we identify with $\mathbb{R}^{n-1}$, and let K_0 be the orthogonal projection of K onto H. Let $f : K_0 \to \mathbb{R}$ be the strictly concave function defining "one half" of $\operatorname{bd} K$. Note that at any point of f, the supporting hyperplane of K is spanned by the vectors $(0, \dots, 0, 1, 0, \dots, 0, \dots, t\partial_i f)$, where the ith coordinate is 1, and $i = 1, 2, \dots, n-1$. Thus, the equilibria correspond to the points where the vector $(x_1, x_2, \dots, x_{n-1}, tf)$ is perpendicular to each of these vectors, and hence to the solutions of the system of equations

$$x_i + t^2 f \partial_i f = 0.$$

From now on we need to repeat the planar argument. □

A more elaborate version of our argument yields the following, stronger statement. To formulate it, let $eq_i(L)$ denote the number of equilibria of index i, of the convex body L. Note that $eq_0(L)$ is the number of stable points, and $eq_n(L)$ is the number of unstable points.

Corollary 2. *$K_0 = H \cap K$. Then, if t is sufficiently large, $eq_n(K(t)) = 2$, and for any $i = 0, 1, \dots, n-1$, $eq_i(K(t)) = eq_i(K_0)$.*

Remark 1. *We note that, following the idea of the proof of Theorem 3, one can obtain statements similar to Theorem 3 and Corollary 2, about the case that $t > 0$ is sufficiently small.*

3. Summary

In this note we studied the evolution of the isoperimetric ratio $I(t)$ under orthogonal affinity. The former is an important geological indicator of abrasion which has been both measured in the field [9] and in laboratory experiments [10], the latter is a candidate model for friction, one important component of abrasion [3]. We also studied the evolution $T(t)$ of the number of static equilibrium points under the same process.

One important conclusion from our results is that under one-parameter orthogonal affinity $I(t)$ and $T(t)$ reach their respective minima simultaneously, as t approaches either zero or infinity. We also showed that $I(t)$ has a single maximum. While the maximum of $T(t)$ often does not coincide with the maximum of $I(t)$, and $T(t)$ may have even several local maxima, we still believe that the global trend of the two functions is related. The one-parameter orthogonal affinity associates with each direction v and each parameter value t a real number $I_v(t) \in [0, 1]$ and an integer $T_v(t)$. One might try to formalize this relation by statistical methods, e.g. by computing the correlation coefficient between $I_v(t)$ and $T_v(t)$, however, correlation is only suitable to detect linear dependence. Instead, we attempt to grasp the relationship by averaging, formulate a conjecture and ask a question related to the averaged system. The latter is defined in the following manner: we subdivide the unit interval into k equal parts $p_i = [i/k, (i+1)/k), i = 0, 1, \dots k-1$.

We denote the average of all $T_v(t)$ such that $I_v(t) \in p_i$ by t_i. More specifically, we let

$$t_i = \frac{\int_0^\infty \frac{1}{\sqrt{1+t^2}} \int_{S_i(t)} T_v(t)\,\mathrm{d}\,v\,\mathrm{d}\,t}{\int_0^\infty \frac{1}{\sqrt{1+t^2}}\,\mathrm{Svol}(S_i(t))\,\mathrm{d}\,t},$$

where $S_i(t) = \{v \in \mathbb{S}^{n-1} : I_v(t) \in p_i\}$, and Svol denotes spherical volume. The largest value of $k = k^\star(K)$ for which the sequence $\{t_i = t_i(K)\}$ is monotonically growing we call the *critical number* associated with the convex body K.

Conjecture 1. *For any convex body K, $k^\star(K) \geq 2$,*

and of particular interest would be a universal lower bound $k_n^\star$ on $k^\star(K)$, depending only on the dimension of K. Needless to say, the higher this lower bound, the stronger it indicates the connection between $I(t)$ and $T(t)$.

References

[1] V.I. Arnold, *Ordinary differential equations*, 10th printing, MIT Press, Cambridge, 1998.
[2] F. Bloore *The shape of pebbles* J. Int. Ass. Math. Geol., 9(2), 113122 (1977).
[3] G. Domokos, G.W. Gibbons, *The evolution of pebble size and shape in space and time* Proc. R.Soc. London A **468**, 3059-3079 (2012).
[4] G. Domokos, G.W. Gibbons, *Geometrical and physical models of abrasion* (2013) arXiv preprint arXiv:1307.5633.
[5] Domokos G., Sipos A.A., Szabo, T., Varkonyi, P.L., *Pebbles, shapes and equilibria*, Math. Geosci. **42** (2010), 29-47.
[6] W. J. Firey, *Isoperimetric ratios of Reuleaux polygons*, Pacific J. Math. **10** (1960), 823-830.
[7] W.J. Firey *The shape of worn stones*, Mathmetika **21**, 1-11 (1974).
[8] M.E. Gage *An isoperimetric inequality with applications to curve shortening*, Duke Math. J. **50** (4), (1983) pp 1225-1229.
[9] K.L. Miller, T. Szabó, D.J. Jerolmack, G. Domokos, *Quantifying the significance of abrasion and selective transport for downstream fluvial grain size evolution*, J. Geophyisocal Research / Earth Surface, in press
[10] F.M. McCubbin et. al. *Alteration of Sedimentary Clasts in Martian Meteorite Northwest Africa 7034* NASA Technical Report JSC-CN-31647, document ID 20140010653.

Gábor Domokos, Dept. of Mechanics, Materials and Structures, Budapest University of Technology, Műegyetem rakpart 1-3., Budapest, Hungary, 1111

E-mail address: domokos@iit.bme.hu

Zsolt Lángi, Dept. of Geometry, Budapest University of Technology and Economics, Budapest, Egry József u. 1., Hungary, 1111

E-mail address: zlangi@math.bme.hu